

Traumatic Spinal Cord Injury—Repair and Regeneration

Christopher S. Ahuja, MD*^{†§||}

Satoshi Nori, MD^{||}

Lindsay Tetreault, MD[§]

Jefferson Wilson, MD*^{§||}

Brian Kwon, MD, PhD*^{**}

James Harrop, MD^{**}

David Choi, MD, PhD^{§§}

Michael G. Fehlings, MD,
PhD*^{†§||}

*Division of Neurosurgery, Department of Surgery, University of Toronto, Toronto, Canada; [†]Institute of Medical Science, University of Toronto, Toronto, Canada; [§]Department of Surgery, University of Toronto, Toronto, Canada; ^{||}Spine Program, University of Toronto, Toronto, Canada; ^{||}Department of Genetics and Development, University of Toronto, Toronto, Canada; [†]Vancouver Spine Institute, Vancouver General Hospital, Vancouver, Canada; ^{**}Department of Surgery, University of British Columbia, Vancouver, Canada; ^{**}Thomas Jefferson University Hospital, Philadelphia, Pennsylvania; ^{§§}National Hospital for Neurology and Neurosurgery, University College London, London, England

Correspondence:

Michael Fehlings, MD, PhD,
Toronto Western Hospital,
West Wing 4th floor 399 Bathurst Street,
Toronto, ON, M5T 2S8
E-mail: Michael.Fehlings@uhn.ca

Received, August 16, 2016.

Accepted, January 12, 2017.

Copyright © 2016 by the
Congress of Neurological Surgeons

BACKGROUND: Traumatic spinal cord injuries (SCI) have devastating consequences for the physical, financial, and psychosocial well-being of patients and their caregivers. Expediently delivering interventions during the early postinjury period can have a tremendous impact on long-term functional recovery.

PATHOPHYSIOLOGY: This is largely due to the unique pathophysiology of SCI where the initial traumatic insult (primary injury) is followed by a progressive secondary injury cascade characterized by ischemia, proapoptotic signaling, and peripheral inflammatory cell infiltration. Over the subsequent hours, release of proinflammatory cytokines and cytotoxic debris (DNA, ATP, reactive oxygen species) cyclically adds to the harsh postinjury microenvironment. As the lesions mature into the chronic phase, regeneration is severely impeded by the development of an astroglial-fibrous scar surrounding coalesced cystic cavities. Addressing these challenges forms the basis of current and upcoming treatments for SCI.

MANAGEMENT: This paper discusses the evidence-based management of a patient with SCI while emphasizing the importance of early definitive care. Key neuroprotective therapies are summarized including surgical decompression, methylprednisolone, and blood pressure augmentation. We then review exciting neuroprotective interventions on the cusp of translation such as Riluzole, Minocycline, magnesium, therapeutic hypothermia, and CSF drainage. We also explore the most promising neuroregenerative strategies in trial today including Cethrin™, anti-NGO antibody, cell-based approaches, and bioengineered biomaterials. Each section provides a working knowledge of the key preclinical and patient trials relevant to clinicians while highlighting the pathophysiologic rationale for the therapies.

CONCLUSION: We conclude with our perspectives on the future of treatment and research in this rapidly evolving field.

KEY WORDS: Spinal cord injury, Trauma, Regenerative medicine, Stem cells, Neuroprotection, Management, Clinical trial

Neurosurgery 80:S9–S22, 2017

DOI:10.1093/neuros/nyw080

www.neurosurgery-online.com


Traumatic spinal cord injuries (SCI) have devastating physical, psychosocial, and vocational implications for patients and caregivers. Direct lifetime costs can reach a staggering \$1.1 to 4.6 million per patient with over 1 million people affected in North America alone (Figure 1).^{1–3} For treating physicians, a working knowledge of current and emerging therapies in SCI is critical to expediently deliver care and improve long-term functional outcomes for patients.^{4,5} This paper summarizes the evidence-based management of a patient with acute SCI and discusses upcoming neuroprotective and neuroregenerative strategies on

the cusp of translation. A primer on the unique pathophysiology of SCI is provided to aid in understanding the rationale behind the diverse range of therapeutic approaches discussed below.

PATHOPHYSIOLOGY

Primary and Secondary SCI

SCI can be categorized into primary and secondary phases.^{6,7} The primary SCI is the result of physical forces of the initial traumatic event and is often the most important determinant of injury severity; physical forces



involved can include compression, shearing, laceration, and acute stretch/distraction.⁸ After the primary injury event, a cascade of secondary injury events is initiated which serves to expand the zone of neural tissue injury and exacerbate neurological deficits and outcomes.^{9,10} Secondary SCI is a delayed and progressive tissue injury following the primary SCI. During this secondary injury cascade, inflammatory cells such as macrophages, microglia, T-cells, and neutrophils infiltrate the injury site as a result of disruption of the blood–spinal cord barrier.¹³⁸ These cells trigger the release of inflammatory cytokines such as tumor necrosis factor- α (TNF- α), interleukin (IL)-1 α , IL-1 β , and IL-6, with tissue levels of these cytokines peaking at 6 to 12 h after injury and remaining elevated up to 4 days after injury.^{11,12} In addition, a loss of ionic homeostasis after SCI results in intracellular hypercalcemia which activates calcium-dependent proteases and causes mitochondrial dysfunction ultimately leading to apoptotic cell death.¹³ Oligodendrocytes are particularly susceptible to apoptotic loss and not just at the site of impact. This apoptotic loss has been observed distant from the epicenter of SCI as well as at the lesion epicenter and leads to demyelination of preserved axons.^{14–16} In addition, phagocytic inflammatory cells release reactive oxygen species which causes DNA oxidative damage, protein oxidation, and lipid peroxidation. Delayed necrosis and apoptosis are further induced by this process.^{17–19} After SCI, upregulated release of excitatory amino acids, such as glutamate and aspartate, is observed due to release from disrupted cells.^{20,21} The excessive activation of excitatory amino acid receptors produces excitotoxicity and


further propagation of loss of neurons and glia by both necrotic and apoptotic cell death.²²

Barriers to Regeneration

It is widely recognized that regeneration of the adult mammalian central nervous system (CNS), including the spinal cord, is difficult due to limited plasticity.²³ Although recent progress in the field of SCI research has demonstrated that the CNS has more inherent regenerative capacity than that was once thought,^{24,25} it does not have the same regenerative capacity that is observed in the peripheral nervous system (PNS). Compared with the PNS, not only is the regenerative capacity of CNS axons lower, but it also decreases with advancing age.²⁶

The inhibitory nature of CNS myelin, which starkly contrasts the effects of PNS myelin, was first recognized in 1985.²⁷ Myelin-associated proteins, including neurite outgrowth inhibitor A (Nogo A),^{28,29} myelin-associated glycoprotein,^{30,31} and oligodendrocyte-myelin glycoprotein,³² bind Nogo receptors to activate the GTPase Rho A. Rho-associated protein kinase (ROCK) is the effector of Rho A, which regulates further downstream effectors, and leads to apoptosis and growth cone collapse of regenerating axons along with neurite retraction.

Additional external barriers potentially add to the inhibition of regeneration. Hypertrophied astrocytes form a physical barrier called the glial scar, which walls off injured tissue from the healthy tissue.³³ The astrocytes also form a chemical barrier by secreting



a number of growth inhibitory chondroitin sulfate proteoglycans (CSPGs) including neurocan, versican, brevican, phosphacan, and NG2.³⁴ Fibroblasts also infiltrate the perilesional region and replace the extracellular matrix with fibrous connective tissue. This is associated with the deposition of inhibitory extracellular matrix molecules which function as chemical barriers to axonal regeneration similar to myelin-associated inhibitors (Figure 2).³⁵

CURRENT MANAGEMENT

The current management of SCI largely follows the American Association of Neurological Surgeons (AANS) and Congress of Neurological Surgeons (CNS) joint section guideline series (Table 1) as well as an upcoming AOSpine 2016 guideline.³⁶ Initial care in the field prioritizes securing the airway, breathing, and circulation followed by early recognition of SCI and rapid referral to specialized centers in order to expedite delivery of time-sensitive interventions.³⁶ To limit further insult to the highly vulnerable cord, spinal immobilization should be performed for all patients with suspected or confirmed injuries.³⁶ This typically involves a rigid cervical collar, backboard for transport, and spinal precautions for patient transfers (eg, logroll maneuver with inline manual cervical stabilization and a transfer board). Systemic hypotension (systolic blood pressure <90 mm Hg), even for brief periods, should be avoided as it is associated with worse long-term neurological outcomes.³⁶ This can be particularly challenging as hypovolemia is common in polytrauma, and interruption of spinal cord sympathetic fibers can induce a profound loss of vascular tone and bradycardia (neurogenic

shock). Resuscitation with large-volume crystalloids is typical; however, alpha-agonists (eg, phenylephrine) or mixed alpha/beta-agonists (eg, dopamine, norepinephrine) may also be required as adjuncts. Once resuscitated, an American Spinal Injury Association (ASIA) International Standards for Neurological Classification of SCI (ISNCSCI) examination should be documented to establish baseline function and the level of neurological injury (Table 2, Figure 3).³⁶

Early localization and classification of osseoligamentous and neurological injuries is necessary to expediently provide the outcome-altering therapies discussed below.^{4,5,37,38} Computerized tomography (CT) imaging is recommended for all patients with suspected SCI as x-rays can miss up to 6% of injuries.³⁹ When evaluating patients with high-energy mechanisms and confirmed cervical injuries, thoracolumbar imaging is recommended to rule out concomitant injuries that may not be clinically apparent.⁴⁰ The role of magnetic resonance imaging (MRI) in the initial workup of patients remains unclear; however, urgent MRI is strongly recommended, particularly in cases with unexplained neurological deficits to rule out ongoing spinal cord compression due to occult ligamentous injuries, epidural hematomas, or critical disk herniations. The utility of MRI in prognostication is also becoming more apparent as validated prediction scores continue to be published.⁴¹

Concurrent with the diagnostic workup, patients should be transferred to a critical care unit providing continuous respiratory, cardiac, and hemodynamic monitoring.³⁶ Immediate life- or limb-threatening injuries should be managed by the appropriate teams while maintaining strict spinal immobilization. Delivering effective care in SCI requires a collaborative

TABLE 1. “Current best practices for the diagnosis and management of SCI. The table displays several key recommendations, many of which are from the 2013 updated guidelines from the Joint Section on Disorders of the Spine and Peripheral Nerves of the American Association of Neurological Surgeons and the Congress of Neurological Surgeons.”^{135,a}

Topic	Level of AANS/CNS recommendation	Guideline/recommendation
Hypotension	Level III	Correction of hypotension to systolic blood pressure > 90 mm Hg) as soon as possible
	Level III	Maintenance of mean arterial blood pressure between 85 and 90 mm Hg for 7 days
Hypoxia	None	Hypoxia (PaO ₂ < 60 mm Hg or O ₂ saturation < 90%) should be avoided [3]
ICU monitoring	Level III	SCI patients should be managed in an intensive care unit (ICU) setting with cardiac, hemodynamic, and respiratory monitoring to detect cardiovascular dysfunction and respiratory insufficiency
Immobilization	Level II	Patients with SCI or suspected SCI (except in penetrating injury) should be immobilized
	Level III	Spinal immobilization should be performed with rigid cervical collar and supportive blocks on a backboard with straps
Specialized centers	Level III	SCI patients should be transferred expediently to specialized centers of SCI care
Examination	Level II	The ASIA ISNCSCI examination should be performed and documented
Imaging	Level I	No cervical imaging is required in awake trauma patients that have no neck pain/tenderness, normal neurological examination, normal range of motion, and no distracting injuries
	Level I	CT is recommended in favor of cervical x-rays
	Level I	CT angiography is recommended in patients that meet the modified Denver screening criteria [9]
	Level I	Methylprednisolone is not recommended ^b
Spinal cord decompression	None	Surgical decompression prior to 24 h after SCI can be performed safely and is associated with improved neurological outcome [10]
	Level III	Early closed reduction of fracture/dislocation in awake patients without a rostral injury is recommended, and pre-reduction MRI does not appear to influence outcome

^aReprinted with permission of Springer from Martin AR, Aleksanderek I, Fehlings MG. Diagnosis and acute management of spinal cord injury: current best practices and emerging therapies. *Current Trauma Reports*. 2015;1(3):169-181.

^bThe authors do not agree with this guideline given the balance of trial data available and the 2012 Cochrane review on steroids for SCI. An upcoming 2016 AOSpine guideline will recommend MPSS as a treatment option for patients within 8 h of injury. See the ‘Steroids for SCI’ section for a full discussion.

multidisciplinary approach including fiberoptic intubation by anesthesia/critical care, modified intraoperative positions for general surgery/orthopedic procedures, and early recognition of therapeutic windows, which can positively alter long-term outcomes.

Early Surgical Decompression

After SCI, ongoing mechanical compression of the spinal cord can impair blood flow causing ischemia and an expanded zone of neural tissue injury. The goal of early surgical decompression after SCI is to relieve this compression, thereby improving the vascular supply to the injured area and limiting the zone of secondary injury expansion. A sizable body of preclinical literature supports the positive effects of early surgical decompression on behavioral and pathological outcomes in animal SCI models.⁴²

With respect to clinical evidence on this topic, a number of comparative cohort studies have been published investigating the clinical impact of performing decompressive surgery prior

to several thresholds. Notably, to investigate the efficacy of early decompression prior to a 24-h threshold, the Surgical Treatment of Acute Spinal Cord Injury Study (STASCIS) enrolled 313 cervical SCI patients.³⁸ Patients receiving early decompression (<24 h after SCI) experienced 2.8 times greater odds of experiencing an ASIA Impairment Scale (AIS) improvement of at least 2 grades at 6 months as compared with patients who underwent late decompression (≥24 h after SCI). Although not statistically significant, early decompression was associated with a reduced incidence of acute in-hospital complications. A prospective Canadian cohort study (including cervical, thoracic, and lumbar SCI, n = 84) confirmed the findings observed in STASCIS, reporting that in the adjusted analysis early decompression was associated with a statistically greater improvement in ASIA motor score recovery at the time of rehabilitation facility discharge.⁴³ Moreover, an observational Canadian cohort study showed that AIS A (complete injury) and AIS B (complete motor injury with incomplete sensory injury) patients who received

TABLE 2. Summary of International Standards for Neurological Classification of Spinal Cord Injury

Parameter	Definition
Motor score	Score out of 100 points representing motor power in 5 key myotomes (each grade out of 5) in each limb
Sensory score	Score out of 224 points representing light touch and pin prick sensation in 28 dermatomes bilaterally
AIS grade	Cumulative measure of injury severity ranging from AIS grade A (most severe motor sensory complete lesion) to AIS grade E (least severe no neurological deficit)
AIS grade A	No motor or sensory preservation below the neurological level of injury (including the distal sacral segments)
AIS grade B	Sensory, but no motor, preservation below the neurological level of injury (including the distal sacral segments)
AIS grade C	Motor preservation below the neurological level of injury (including the distal segments) with less than half of key muscles below the neurological level graded antigravity or better
AIS grade D	Motor preservation below the neurological level of injury (including the distal segments) with at least half of key muscles below the neurological level graded antigravity or better
AIS grade E	Neurological normal in a patients who previously had deficit
Neurological level of injury	The lowest segment where motor and sensory function is normal on both sides
Zone of partial preservation	In AIS grade A patient, lowest dermatome or myotome with partial innervation

Based on Kirshblum et al. J Spinal Cord Med. 2011;34(6):535-546.¹³⁶

early decompression experienced shorter length of hospital stay, while AIS B, C, and D incomplete injury patients decompressed in an early fashion demonstrated an additional 6.3 points in motor recovery as compared to those decompressed late.⁴ Taken together, these findings support the concept of “Time is Spine”, emphasizing the importance of early diagnosis and intervention to enhance long-term outcomes.

Central cord injury is the most common form of incomplete SCI. It is defined as greater weakness in the upper extremities than the lower extremities, typically present without spinal instability, with patients typically experiencing substantial spontaneous neurological recovery. Historically, early decompression has been avoided in cases of central cord injury due findings of poor outcomes after surgery.⁴⁴ However, an analysis of prospective data performed by the Spine Trauma Study Group associated early decompression (<24 h after SCI) with an additional 6.3 points of ASIA motor score recovery and 2.8 times odds of AIS grade improvement at 12-month follow-up as compared to late decompression (≥24 h after SCI).⁴⁵ In 2013, a randomized control trial Comparing Surgical Decompression Versus Conservative Treatment in Incomplete Spinal Cord Injury (NCT01367405; n = 72) trial was initiated by Raboud University. The study will compare surgical decompression within 24 h to normal conservative treatment without surgery and is currently recruiting participants.

Steroids for SCI

Methylprednisolone (MPSS) is a potent synthetic glucocorticoid which upregulates anti-inflammatory cytokine release

and reduces oxidative stress to enhance neural cell survival in preclinical models of traumatic SCI. The National Acute Spinal Cord Injury Study trial series (1990,⁴⁶ 1997⁴⁷) found an increase in the number of infection-related complication (eg, severe sepsis, severe pneumonia) with the high-dose 48-h protocol, which outweighed the potential neurological benefits. However, a shorter 24-h course of IV MPSS (30 mg/kg bolus + 5.4 mg/kg/h × 23 h) had a substantially lower complication rate and, when administered to a subgroup of patients within 8 h of injury, was still found to improve neurological outcomes. These subgroup analyses and the purported methodology have been a source of controversy for the last 3 decades. A 2012 Cochrane review meta-analysis pooling 6 key randomized trials and observational studies was completed which found that patients receiving MPSS within 8 h of injury had a 4-point greater ASIA motor score improvement.⁴⁸ This modest benefit can have tremendous functional implications for patients when those motor points are recovered in key myotomes such as hand strength and deltoid function. As a result, an upcoming AOSpine 2016 guideline developed by an international expert panel will suggest 24 h of IV MPSS be considered within 8 h of injury for patients without significant medical contraindication.

Blood Pressure Augmentation

Vascular injury and localized edema contribute to ongoing ischemia in the perilesional region. Blood pressure augmentation has emerged as a viable strategy to neuroprotect at-risk tissue by enhancing perfusion. Current AANS/CNS guidelines recommend maintenance of mean arterial pressure

ASIA INTERNATIONAL STANDARDS FOR NEUROLOGICAL CLASSIFICATION OF SPINAL CORD INJURY (ISNCSCI) **ISCS**

Patient Name _____ Date/Time of Exam _____
 Examiner Name _____ Signature _____

RIGHT

MOTOR KEY MUSCLES

Elbow flexors **C5**

Wrist extensors **C6**

Elbow extensors **C7**

Finger flexors **C8**

Finger abductor (little finger) **T1**

UER (Upper Extremity Right)

LER (Lower Extremity Right)

Hip flexors **L2**

Knee extensors **L3**

Ankle dorsiflexors **L4**

Long toe extensors **L5**

Ankle plantar flexors **S1**

(VAC) Voluntary Anal Contraction (Yes/No) ☐

RIGHT TOTALS (MAXIMUM) (50) (56) (56)

MOTOR SUBSCORES

UER ☐ + **UEL** ☐ = **UEMS TOTAL** ☐ (MAX(25) (25) (50))

LER ☐ + **LEL** ☐ = **LEMS TOTAL** ☐ (MAX(25) (25) (50))

NEUROLOGICAL LEVELS

1. SENSORY **R** ☐ **L** ☐

2. MOTOR **R** ☐ **L** ☐

3. NEUROLOGICAL LEVEL OF INJURY (NLI) ☐

4. COMPLETE OR INCOMPLETE ☐

Incomplete = Any sensory or motor function in S4-5

5. ASIA IMPAIRMENT SCALE (AIS) ☐

(In complete injuries only)

ZONE OF PARTIAL PRESERVATION Most caudal level with any innervation

SENSORY **R** ☐ **L** ☐

MOTOR **R** ☐ **L** ☐

SENSORY KEY SENSORY POINTS

Light Touch (LTR) Pin Prick (PPR)

LEFT

MOTOR KEY MUSCLES

C5 Elbow flexors

C6 Wrist extensors

C7 Elbow extensors

C8 Finger flexors

T1 Finger abductor (little finger)

UEL (Upper Extremity Left)

LEL (Lower Extremity Left)

Hip flexors **L2**

Knee extensors **L3**

Ankle dorsiflexors **L4**

Long toe extensors **L5**

Ankle plantar flexors **S1**

(DAP) Deep Anal Pressure (Yes/No) ☐

LEFT TOTALS (MAXIMUM) (50) (56) (56)

MOTOR SUBSCORES

LTR ☐ + **LTL** ☐ = **LT TOTAL** ☐ (MAX(56) (56) (112))

PPR ☐ + **PPL** ☐ = **PP TOTAL** ☐ (MAX(56) (56) (112))

This form may be copied freely but should not be altered without permission from the American Spinal Injury Association. REV 11/15

FIGURE 3. International Standards for Neurological Classification of Spinal Cord Injury clinical examination form. The standardized assessment and calculation of motor and sensory scores is demonstrated on this template. Reprint of the 2015 American Spinal Injury Association and International Spinal Cord Society ISNCSCI assessment form retrieved from http://asia-spinalinjury.org/wp-content/uploads/2016/02/International_Stds_Diagram_Worksheet.pdf.

(MAP) ≥ 85 to 90 mm Hg for 7 days postinjury as this has been found to enhance long-term AIS grade outcomes for patients.⁵ In application, this most often necessitates invasive blood pressure monitoring, IV fluid therapy, and central venous access for continuous infusion of vasopressors.⁴⁹ These substantial requirements have prompted a noninferiority trial entitled Mean Arterial blood Pressure Treatment for Acute Spinal Cord Injury (NCT02232165) comparing MAP ≥ 85 mm Hg vs MAP ≥ 65 mm Hg with results expected by 2017.³

These requirements can also be a significant hindrance to early mobilization, an important component of cardiorespiratory and dermatologic complication prevention. A collaborative interdisciplinary approach utilizing adjunctive measures such as prophylactic vasopressors, abdominal binding, and assistive devices is often required to safely elevate patients. The precise timing of

mobilization is dictated by the patient's hemodynamic status and the expertise of the treating team.⁵⁰

KEY TRIALS IN NEUROPROTECTION

In addition to early decompression, MAP augmentation and IV MPSS, several other neuroprotective strategies targeting key components of the secondary injury cascade have emerged in preclinical research.⁵¹ The most promising therapies currently being translated are discussed in this section.

Pharmacological Therapies

Riluzole

Riluzole is a benzothiazole sodium channel blocker currently approved by the US Food and Drug Administration, European

Medicines Agency, and Health Canada for the treatment of amyotrophic lateral sclerosis.^{52,53} It protects against excitotoxic cell death by blocking sodium influx in injured neurons and restricting the presynaptic release of glutamate.⁵⁴ Animal studies in SCI have demonstrated its ability to reduce neuronal loss and cavity size while improving sensorimotor and electrophysiological outcomes.⁵⁵⁻⁵⁸ A collaborative effort to study Riluzole for SCI is being led by the corresponding author (MGF) and includes the North American Clinical Trials Network, AOSpine, the Ontario Neurotrauma Foundation, and the Rick Hansen Institute. This phase II/III randomized controlled trial (RCT) (n = 351) entitled “Riluzole in Spinal Cord Injury Study” (RISCIS; NCT01597518) is currently recruiting patients with acute C4-8 ASIA grade A/B/C injuries and will assess multiple outcomes including the AIS, Brief Pain Inventory, and Spinal Cord Independence Measure.³ The study is expected to conclude in 2018.

Magnesium

Magnesium can act as an N-methyl-D-aspartate (NMDA) receptor antagonist to decrease excitotoxicity and also functions as an anti-inflammatory agent. Stable cerebrospinal fluid (CSF) levels can be generated by delivering magnesium with an excipient such as polyethylene glycol (PEG).⁵⁹⁻⁶¹ In animal models, the Mg-PEG combination has been shown to enhance tissue sparing and lead to behavioral recovery.^{62,63} A phase I trial (n = 15; NCT01750684) of an Mg-PEG combination (AC105) led by Acorda Therapeutics Inc. (Ardsley, New York) concluded in February 2015 with results pending report.³

Minocycline

Minocycline is a second-generation bacteriostatic tetracycline antibiotic that has demonstrated neuroprotective properties in preclinical models of CNS disorders including Huntington's disease and multiple sclerosis.^{64,65} This stems in part from its significant anti-inflammatory effect mediated by inhibition of microglial activation, IL-1 β , TNF- α , cyclooxygenase-2, and matrix metalloproteinases.⁶⁶⁻⁶⁹ In animal studies, minocycline treatment after acute SCI has been shown to reduce lesion size and promote tissue sparing.^{70,71} A phase II trial demonstrated that patients with acute incomplete cervical SCI (n = 25) may benefit from early minocycline administration as they found a 14-point ASIA motor score improvement compared to placebo (P = .05).⁷² This exciting result led to the development of a phase III trial (n = 248) entitled ‘Minocycline in Acute Spinal Cord Injury’ (NCT01828203) which will assess intravenously administered minocycline for 7 days vs placebo and is expected to report by 2018.³

Monosialotetrahexosylganglioside (GM-1) Ganglioside

GM-1 is a glycosphingolipid found in cell membranes with the ability to activate receptor tyrosine kinases to enhance neural plasticity and regeneration. It has been successfully used for neuroprotection in animal models of SCI where it enhanced tissue

sparing.⁷³ A successful phase II trial (n = 37) found improved 1-year ASIA motor scores for those receiving daily GM-1 for 18 to 32 days postinjury.⁷⁴ Unfortunately, a follow-up phase III RCT (n = 797) found no statistically significant improvement with treatment.⁷⁵ No further studies have been registered.

Fibroblast Growth Factor

Fibroblast growth factor is a heparin-binding protein found to be neuroprotective against excitotoxicity while also reducing oxygen-free radical generation.⁷⁶ In animal models, it has been shown to reduce motor neuron loss and improve respiratory deficits.^{77,78} A phase I/II trial (n = 62; NCT01502631) of the fibroblast growth factor-analog, SUN13837 (Asubio Pharmaceuticals Inc., Edison, New Jersey), completed in 2015 with results pending publication.³

Granulocyte Colony-stimulating Factor

Granulocyte colony-stimulating factor (G-CSF; CSF 3) is a cytokine glycoprotein found in numerous tissues throughout the body. It is capable of promoting cell proliferation, survival, and mobilization. In the CNS, it has been shown to facilitate survival of ischemic cells and reduce inflammatory cytokine expression (eg, TNF- α , IL-1 β).⁷⁹⁻⁸¹ A recent pair of nonrandomized phase I/IIa trials showed no increase in serious adverse events with G-CSF administration while also demonstrating improvement in AIS outcomes.^{82,83} Additional well-designed RCTs will be required to establish the efficacy of G-CSF for SCI.


Hepatocyte Growth Factor

Hepatocyte growth factor (HGF) is a prosurvival, promotility c-Met receptor ligand. In small animal SCI models, HGF increases neuron survival and decreases oligodendrocyte apoptosis resulting in improved behavioral outcomes.⁸⁴⁻⁸⁶ More recently, HGF has been shown to promote angiogenesis and enhance upper limb recovery in a primate model of cervical SCI.⁸⁵ A phase I/II randomized trial (n = 48; NCT02193334) of KP-100IT (HGF; Kringle Pharma Inc., Osaka, Japan) is now underway with results expected in 2017.³

Nonpharmacologic Therapies

Therapeutic Hypothermia

Therapeutic hypothermia (TH; 32°C-34°C) significantly reduces the basal metabolic rate of the CNS and decreases inflammatory cell activation.⁸⁷ It has been successfully applied in neonatal hypoxic-ischemic encephalopathy and after in-hospital cardiac arrest.⁸⁸⁻⁹⁰ In preclinical SCI models, it has been shown to enhance tissue sparing and promote behavioral recovery prompting a pilot study (n = 14) of early systemic TH for patient with AIS A injuries which found no increase in complication rates and a trend toward increased neurological recovery (43% vs 21%).^{91,92} A phase II/III trial entitled Acute Rapid Cooling Therapy for Injuries of the Spinal Cord has been planned to definitively assess efficacy.



CSF Drainage

CSF drainage attempts to prevent cord hypoperfusion in the critical postinjury period by relieving pressure analogous to external ventricular drainage (EVD) for raised intracranial pressure (ICP). A phase I/II trial ($n = 22$) completed in 2009 found no significant improvement outcomes with drainage; however, the study was not sufficiently powered to demonstrate efficacy.⁹³ Recent large-animal trials have found that CSF drainage acts synergistically with MAP augmentation to improve cord blood flow.⁹⁴ Based on these key results, a phase IIB trial ($N = 60$; NCT02495545) of MAP elevation with CSF drainage has been launched with results expected by December 2017.³

KEY TRIALS IN NEUROREGENERATION

While timely neuroprotective interventions can have tremendous benefits in the acute injury period, the majority of our patients are in the chronic phase of their injuries where

further recovery is limited. This section discusses emerging neuroregenerative therapies in clinical trial or on the cusp of translation (illustrated in Figure 4).

Pharmacological Therapies

Rho-ROCK Inhibitor

Components of the injured adult CNS including CSPGs, myelin-associated glycoproteins, and Nogo potentially inhibit axon outgrowth and attempts at regeneration via the Rho-ROCK signaling pathway. Cethrin/VX-210 (Vertex Pharmaceuticals, Boston, Massachusetts) is a direct Rho inhibitor applied intraoperatively within a fibrin glue sealant to the epidural space.⁹⁵ A mixed open-label phase I/IIa trial ($n = 48$; NCT00500812) of patients with cervical or thoracic injuries found no increase in serious adverse events and a significant improvement in long-term motor scores (18.5 ASIA points) for cervical patients.⁹⁶ These very exciting results have led to a phase III trial in patients with acute cervical SCI which has commenced in 2016.

Anti-NGO Antibody

Anti-NGO is a monoclonal antibody against NGO-A, a major inhibitor component of adult CNS myelin. Anti-NGO treatment delivered by intrathecal injection has been shown to promote axonal sprouting and functional recovery in animal models by clearing the source of this inhibitory signaling.⁹⁷ A phase I trial (n = 51; NCT004060160) of humanized anti-NGO antibody (ATI-355; Novartis, Basel, Switzerland) has been completed with results pending dissemination.³ A European group led by Professor Armin Curt is planning on carrying forward with additional trials with the anti-NGO antibody in patients with SCI.

Nonpharmacologic Therapies

Spinal Cord Stimulation

Spinal cord stimulation (SCS) has been used to successfully treat refractory chronic pain for disorders ranging from SCI to phantom limb pain. In animal models, weakly imposed DC electrical fields have been shown to promote cathode-directed neurite outgrowth while oscillating DC fields can promote bidirectional regeneration.⁹⁸ Recently, a small human study has shown that SCS combined with rehabilitation can provide gradual recovery of voluntary lower limb movement even years after a complete injury.⁹⁹ Additional trials (NCT02592668, NCT02313194) are now ongoing to assess safety/feasibility and validate this exciting finding with results expected by 2018. Another potential application, shown in a rodent model, is the application of SCS in combination with stem cell transplants to enhance the survival and migration (galvanotaxis) of grafted cells beyond the region of injection.

Cell Therapies

Cell-based regenerative therapies are an exciting field as transplanted cells are capable of filling many roles including providing trophic support, modulating the inflammatory response, regenerating lost neural circuits, and remyelinating denuded axons.¹⁰⁰⁻¹⁰² Early research utilized embryonic stem cells (ESCs), however, ethical concerns and limited supplies have driven the field towards induced pluripotent stem cell (iPSCs) which can be derived from any somatic cell, including autologous sources.¹⁰³ While unanticipated challenges have arisen, including early senescence and retained epigenetic memory, iPSCs remain a key therapeutic approach moving forward. Numerous animal studies over the last 3 decades have demonstrated the beneficial effects of a range of transplanted cell types. The most clinically relevant approaches are discussed here.

Neural Stem/Precursor Cells

Derived from stem cells, multipotent neural precursor cells (NPCs) are capable of differentiating to CNS-specific neurons, oligodendrocytes, and astrocytes, making them a particularly promising strategy. In animal studies, they are capable of integrating with host circuits to enhance behavioral recovery

over several weeks.^{104,105} A pair of phase II trials by Stem Cells Inc. (Newark, California) of human CNS stem cell transplants for cervical (n = 31; NCT02163876) and thoracic (n = 12; NCT01321333) injury were terminated in 2016 prior to completion. While results regarding sensorimotor outcomes are pending dissemination, preliminary data suggest no increase in complications rates related to the treatments.³ This provides confirmation of existing safety data that intraparenchymal stem cell transplants are feasible and suggests further optimization of the cells and/or their environment is required to produce meaningful changes in functional recovery.

A parallel strategy to specifically target postinjury demyelination is the transplant of oligodendrocyte precursor cells (OPCs) which preferentially differentiate to functional oligodendrocytes. Asterias Biotherapeutics Inc. (Fremont, California) has launched an open-label phase I/II dose-escalation trial (n = 35; NCT02302157) of their AST-OPC1 cell line with long-term outcome measures including adverse events and serial ISNCSCI exams. The study is expected to complete in 2018.³

Mesenchymal Stem Cells

Mesenchymal stem cells (MSCs) are multipotent cells capable of repairing connective tissues by differentiating to myocytes, osteoblasts, chondrocytes, and adipocytes.¹⁰⁶ They can also modulate local and systemic inflammation which has been exploited in animal models of SCI where MSC treatment led to a decrease in peripheral inflammatory cell infiltration and an increase in parenchymal tissue volume.¹⁰⁷⁻¹¹¹ A phase II/III RCT (n = 32; NCT01676441) by Pharmicell Co. (Seoul, South Korea) studying intraparenchymal and intrathecal MSC treatment for patients with acute AIS B injuries is ongoing with results expected in 2016.³

Schwann Cells

The robust regeneration seen in the PNS is thought to be mediated in large part by Schwann cells (SCs). In animal models of SCI, peripheral SCs transplanted into the CNS were found to remyelinate axons, reduce cystic cavitation, and enhance recovery.¹¹² An open-label phase I trial (n = 10; NCT02354625) by the Miami Project to Cure Paralysis is now investigating SCs in the treatment of patients with chronic AIS A, B, and C cervical or thoracic injuries with results expected by 2018.³

Olfactory Ensheathing Cells

Olfactory ensheathing cells (OECs) protect olfactory neurons exposed to the harsh conditions of the nasal mucosa. They rapidly phagocytose debris and microbes while also providing trophic support through growth factor signaling.¹¹³⁻¹¹⁶ In animal models, OECs have been shown to induce regeneration in reapposed dorsal roots after brachial plexus avulsion injuries resulting in substantial recovery of proprioception.¹¹⁷ Clinical studies of OECs for brachial plexus avulsions are now in preparation by researchers in the UK. OECs harvested from the

nasal mucosa have also been transplanted into the spinal cord for SCI and have been shown to improve neurite outgrowth and endogenous remyelination resulting in impressive behavioral recovery in animal models.¹¹⁸ Numerous clinical trials of OECs for chronic SCI have been completed worldwide and analyzed in a meta-analysis (cumulative $n = 1193$) which found no significant increase in complication rates related to the transplant. Efficacy could not be definitively established due to the quality of the studies.¹¹⁹

Biomaterials

Regeneration is often hindered by the presence of a substantial postinjury cystic cavity, which lacks the substrate to support cell migration and axon growth. Biomaterials have emerged as an exciting strategy to fill cavitation defects and reproduce the complex structural architecture of the extracellular matrix.¹²⁰⁻¹²⁴ Many of these materials can be engineered to biodegrade over time, release growth factors, and can even be seeded with stem cells to enhance engraftment.^{97,98} Several biomaterials have been shown to be effective in animal models of SCI from the acute to chronic phases (eg, HAMC,¹²¹ QL6,^{125,126} fibrinogen,¹²⁷ etc). As the technology evolves, more niche-specific biomaterials are expected to emerge with extended drug-release and cell support capabilities. Currently, a phase III trial ($n = 20$; NCT02138110) of InVivo Therapeutics' Neuro-Spinal Scaffold (Cambridge, Massachusetts) is now recruiting patients with acute AIS A thoracic injuries.³ Results are expected in 2017.

FUTURE DIRECTIONS

The next substantial changes in the management of patients with SCI are likely to be translated from research that adapts to the heterogeneity of SCI. Modified trial designs which specifically target SCI subpopulations are likely to have the greatest impacts on long-term functional recovery. Stratifying patients in this way will require a combination of existing metrics (eg, clinical exam, radiography) and novel assessment techniques (eg, advanced imaging, biochemical biomarkers).

While many novel treatments show promise in animal models of SCI, these experimental paradigms typically involve very controlled injury and recovery conditions after biomechanically precise injuries in animals matched for age, weight, gender, species, and, in some cases, genetic background. This obviously pales in comparison to the natural variability that occurs in the acute human SCI setting. The appreciation of the heterogeneity of human SCI is partly the result of the challenges that have been experienced in the execution of clinical trials of novel therapies, particularly in the acute setting. Variability in neurological recovery requires that many patients be recruited to complete such acute clinical trials in order to be sufficiently powered. Difficulties in achieving such recruitment has plagued the conduct of virtually all acute clinical studies, and the failure to enroll sufficiently large patient cohorts within realistic time

frames has resulted in the premature cessation of numerous clinical trial programs. New approaches to overcome this will be needed in the future to facilitate the conduct of such clinical trials and enhance the speed with which novel treatments for SCI can be validated. Such approaches include narrowing the inclusion window to be more specific in the types and severities of cord injuries being studied, and establishing objective biomarkers for the stratification of injury severity and more precise prediction of neurological outcome.

Seminal large-scale clinical trials for SCI have typically used broad inclusion criteria to bolster recruitment across participating centers. However, post hoc subgroup analyses have now demonstrated that patient characteristics, presentations, and the underlying pathophysiology in SCI can be highly heterogeneous which can influence the relationship between treatments and outcomes.^{46-48,128,129} As a result, more recent studies are recruiting carefully selected populations. The upcoming Riluzole in Spinal Cord Injury (RISCIS; NCT01597518) trial is an example where recruitment is limited to patients with C4-8 injuries and ASIA grades A, B, or C.^{3,58} Other clinical initiatives have similarly restricted inclusion both with regard to the level of injury (cervical vs thoracic), severity of injury (AIS grade A, B, or C), and timing of intervention. While logistically demanding, this careful selection will allow a more valid assessment of the drug's efficacy.

The next generation of trials will also need to further define subpopulations based on quantifiable imaging and biochemical biomarkers. MRI is a key imaging modality for most CNS pathologies; however, its adoption in SCI trials has been limited. This is likely because the most common sequences (T1- and T2-weighted) rely on gross measurements of hemorrhage and compression providing only modest utility in predicting outcomes. Future MR imaging will need to quantify the cord microstructure to better estimate damage and recovery potential. Emerging techniques for this include diffusion tensor imaging (axon integrity), myelin water fraction (myelination), MR spectroscopy (gliosis or ischemia), and functional MRI (connectivity).^{130,131}

Biochemical biomarkers are also being extensively explored. The Canadian Multicentre CSF Monitoring and Biomarker Study (NCT01279811) is testing CSF over 5 days for inflammatory cell proteins, interleukins, and other cytokines.³ Specific proteins, such as IL-6, S100 β , and tau within the CSF of acute SCI patients have been shown to be able to objectively stratify injury severity and predict AIS grade and motor score improvement.^{132,133} An additional class of biomarkers currently under study through the Rick Hansen Institute is micro RNAs (miRNA), which are short noncoding RNA segments that can regulate post-transcriptional gene expression. miRNAs are specifically up- or downregulated with varying grades of SCI, and may hold important prognostic information as they are further understood.¹³⁴ Together these biomarkers will yield important data to help identify subgroups within the heterogeneous SCI population, and when combined with clinical examination, will

allow patients to be stratified by their specific pathophysiologic niche into targeted trials.

CONCLUSION

The breadth of therapeutic approaches discussed within this review and the rapidly evolving management of a patient with SCI highlight the excitement and progress continuing to be made in the field. The collaborative effort of thousands of physicians, scientists, and allied health professionals has generated a treatment pipeline with numerous promising therapies likely to see translation over the next decade.

Disclosures

This work was funded by funded by AOSpine North America. The authors have no personal, financial, or institutional interest in any of the drugs, materials, or devices described in this article.

REFERENCES

- Center NSCIS. Spinal cord injury facts and figures at a glance. *J Spinal Cord Med.* 2014;37(1):117-118.
- Foundation CaDR. *One Degree of Separation: Paralysis and Spinal Cord Injury in the United States*, 2010. Available at: http://www.christopherreeve.org/site/c.ddJFKRNoFiG/b.5091685/k.58BD/One_Degree_of_Separation.htm. Accessed September 10, 2016.
- ClinicalTrials.gov. 2016. <https://clinicaltrials.gov/>. Accessed August 4, 2016.
- Dvorak MF, Fallah N, Fisher CG, et al. The influence of time from injury to surgery on motor recovery and length of hospital stay in acute traumatic spinal cord injury: an observational Canadian cohort study. *J Neurotrauma.* 2015;32(9):645-654.
- Wilson JR, Forgione N, Fehlings MG. Emerging therapies for acute traumatic spinal cord injury. *CMAJ.* 2013;185(6):485-492.
- Tator CH. Update on the pathophysiology and pathology of acute spinal cord injury. *Brain Pathol.* 1995;5(4):407-413.
- McDonald JW, Sadowsky C. Spinal-cord injury. *Lancet.* 2002;359(9304):417-425.
- Ackery A, Tator C, Krassioukov A. A global perspective on spinal cord injury epidemiology. *J Neurotrauma.* 2004;21(10):1355-1370.
- Yip PK, Malaspina A. Spinal cord trauma and the molecular point of no return. *Mol Neurodegener.* 2012;7:6.
- Norenberg MD, Smith J, Marcillo A. The pathology of human spinal cord injury: defining the problems. *J Neurotrauma.* 2004;21(4):429-440.
- Nakamura M, Houghtling RA, MacArthur L, Bayer BM, Bregman BS. Differences in cytokine gene expression profile between acute and secondary injury in adult rat spinal cord. *Exp Neurol.* 2003;184(1):313-325.
- Uldredj A, Chio JC, Ahuja CS, Fehlings MG. Modulating the immune response in spinal cord injury. *Expert Rev Neurother.* 2016;16(10):1-3.
- Schanne FA, Kane AB, Young EE, Farber JL. Calcium dependence of toxic cell death: a final common pathway. *Science.* 1979;206(4419):700-702.
- Beattie MS, Farooqui AA, Bresnahan JC. Review of current evidence for apoptosis after spinal cord injury. *J Neurotrauma.* 2000;17(10):915-925.
- Crowe MJ, Bresnahan JC, Shuman SL, Masters JN, Beattie MS. Apoptosis and delayed degeneration after spinal cord injury in rats and monkeys. *Nat Med.* 1997;3(1):73-76.
- Dong H, Fazzaro A, Xiang C, Korsmeyer SJ, Jacquin MF, McDonald JW. Enhanced oligodendrocyte survival after spinal cord injury in Bax-deficient mice and mice with delayed Wallerian degeneration. *J Neurosci.* 2003;23(25):8682-8691.
- Waxman SG. Demyelination in spinal cord injury. *J Neuro Sci.* 1989;91(1-2):1-14.
- Dizdaroğlu M, Jaruga P, Birincioglu M, Rodriguez H. Free radical-induced damage to DNA: mechanisms and measurement. *Free Radic Biol Med.* 2002;32(11):1102-1115.
- Hausmann ON. Post-traumatic inflammation following spinal cord injury. *Spinal Cord.* 2003;41(7):369-378.
- Liu M, Wu W, Li H, et al. Necroptosis, a novel type of programmed cell death, contributes to early neural cells damage after spinal cord injury in adult mice. *J Spinal Cord Med.* 2015;38(6):745-753.
- Wang Y, Wang H, Tao Y, Zhang S, Wang J, Feng X. Necroptosis inhibitor necrostatin-1 promotes cell protection and physiological function in traumatic spinal cord injury. *Neuroscience.* 2014;266:91-101.
- Li S, Stys PK. Mechanisms of ionotropic glutamate receptor-mediated excitotoxicity in isolated spinal cord white matter. *J Neurosci.* 2000;20(3):1190-1198.
- Cajal RY. *Degeneration and Regeneration of the Nervous System*. London: Oxford University Press; 1928.
- Barnabe-Heider F, Frisen J. Stem cells for spinal cord repair. *Cell Stem Cell.* 2008;3(1):16-24.
- Okano H, Yamanaka S. iPS cell technologies: significance and applications to CNS regeneration and disease. *Mol Brain.* 2014;7:22.
- Kwon BK, Liu J, Messerer C, et al. Survival and regeneration of rubrospinal neurons 1 year after spinal cord injury. *Proc Natl Acad Sci USA* 2002;99(5):3246-3251.
- Schwab ME, Thoenen H. Dissociated neurons regenerate into sciatic but not optic nerve explants in culture irrespective of neurotrophic factors. *J Neurosci.* 1985;5(9):2415-2423.
- Chen MS, Huber AB, van der Haar ME, et al. Nogo-A is a myelin-associated neurite outgrowth inhibitor and an antigen for monoclonal antibody IN-1. *Nature.* 2000;403(6768):434-439.
- Freund P, Schmidlin E, Wannier T, et al. Nogo-A-specific antibody treatment enhances sprouting and functional recovery after cervical lesion in adult primates. *Nat Med.* 2006;12(7):790-792.
- Cafferty WB, Duffy P, Huebner E, Strittmatter SM. MAG and OMgp synergize with Nogo-A to restrict axonal growth and neurological recovery after spinal cord trauma. *J Neurosci.* 2010;30(20):6825-6837.
- DeBellard ME, Tang S, Mukhopadhyay G, Shen YJ, Filbin MT. Myelin-associated glycoprotein inhibits axonal regeneration from a variety of neurons via interaction with a sialoglycoprotein. *Mol Cell Neurosci.* 1996;7(2):89-101.
- Barton WA, Liu BP, Tzvetkova D, et al. Structure and axon outgrowth inhibitor binding of the Nogo-66 receptor and related proteins. *EMBO J.* 2003;22(13):3291-3302.
- Guha A, Tator CH, Rochon J. Spinal cord blood flow and systemic blood pressure after experimental spinal cord injury in rats. *Stroke.* 1989;20(3):372-377.
- McKeon RJ, Schreiber RC, Rudge JS, Silver J. Reduction of neurite outgrowth in a model of glial scarring following CNS injury is correlated with the expression of inhibitory molecules on reactive astrocytes. *J Neurosci.* 1991;11(11):3398-3411.
- Brazda N, Muller HW. Pharmacological modification of the extracellular matrix to promote regeneration of the injured brain and spinal cord. *Prog Brain Res.* 2009;175:269-281.
- Resnick DK. Updated guidelines for the management of acute cervical spine and spinal cord injury. *Neurosurgery.* 2013;72(suppl 2):1.
- Wilson JS, Craven AC, Verrier M, et al. Early versus late surgery for traumatic spinal cord injury: the results of a prospective Canadian cohort study. *Spinal Cord.* 2012;50(11):840-843.
- Fehlings MG, Vaccaro A, Wilson JR, et al. Early versus delayed decompression for traumatic cervical spinal cord injury: results of the Surgical Timing in Acute Spinal Cord Injury Study (STASCIS). *PLoS One.* 2012;7(2):e32037.
- Ryken TC, Hadley MN, Walters BC, et al. Radiographic assessment. *Neurosurgery.* 2013;72(suppl 2):54-72.
- Sixta S, Moore FO, Dittillo MF, et al. Screening for thoracolumbar spinal injuries in blunt trauma: an Eastern Association for the Surgery of Trauma practice management guideline. *J Trauma Acute Care Surg.* 2012;73(5 suppl 4):S326-S332.
- Wilson JR, Grossman RG, Frankowski RF, et al. A clinical prediction model for long-term functional outcome after traumatic spinal cord injury based on acute clinical and imaging factors. *J Neurotrauma.* 2012;29(13):2263-2271.

42. Batchelor PE, Wills TE, Skeers P, et al. Meta-analysis of pre-clinical studies of early decompression in acute spinal cord injury: a battle of time and pressure. *PLoS One*. 2013;8(8):e72659.
43. Wilson JR, Singh A, Craven C, et al. Early versus late surgery for traumatic spinal cord injury: the results of a prospective Canadian cohort study. *Spinal Cord*. 2012;50(11):840-843.
44. Schneider RC, Cherry G, Pantek H. The syndrome of acute central cervical spinal cord injury; with special reference to the mechanisms involved in hyperextension injuries of cervical spine. *J Neurosurg*. 1954;11(6):546-577.
45. Lenehan B, Fisher CG, Vaccaro A, Fehlings M, Aarabi B, Dvorak MF. The urgency of surgical decompression in acute central cord injuries with spondylosis and without instability. *Spine*. 2010;35(21 suppl):S180-S186.
46. Bracken MB, Shepard MJ, Collins WF, et al. A randomized, controlled trial of methylprednisolone or naloxone in the treatment of acute spinal-cord injury. Results of the Second National Acute Spinal Cord Injury Study. *N Engl J Med*. 1990;322(20):1405-1411.
47. Bracken MB, Shepard MJ, Holford TR, et al. Administration of methylprednisolone for 24 or 48 hours or tirilazad mesylate for 48 hours in the treatment of acute spinal cord injury. Results of the Third National Acute Spinal Cord Injury Randomized Controlled Trial. National Acute Spinal Cord Injury Study. *JAMA*. 1997;277(20):1597-1604.
48. Bracken M. Steroids for acute spinal cord injury. *Cochrane Database of Systematic Reviews*. Jan 2012.
49. Ahuja C, Fehlings MG. Spinal cord injury. In: Reich D, Mayer S, Uysal S, eds. *Neuroprotection in Critical Care and Perioperative Medicine*. New York, NY: Oxford University Press; 2017.
50. Ahuja CM, Martin AR, Fehlings M. Recent advances in managing a spinal cord injury secondary to trauma. *F1000 Faculty Reviews*. 2016;5:1017.
51. Ahuja C, Fehlings M. Neuroprotection of the injured spinal cord – what does the future hold? In: Bellabarba C, Kandziora F, eds, Vialle LRG(series ed). *AO Spine Master Series*. Vol 7: 89-106. New York, NY: Thieme; 2016.
52. Bhatt JM, Gordon PH. Current clinical trials in amyotrophic lateral sclerosis. *Exp Opin Invest Drugs*. 2007;16(8):1197-1207.
53. Bensimon G, Lacomblez L, Delumeau JC, Bejuit R, Truffinet P, Meininger V. A study of riluzole in the treatment of advanced stage or elderly patients with amyotrophic lateral sclerosis. *J Neurol*. 2002;249(5):609-615.
54. Azbill RD, Mu X, Springer JE. Riluzole increases high-affinity glutamate uptake in rat spinal cord synaptosomes. *Brain Res*. 2000;871(2):175-180.
55. Nogradi A, Szabo A, Pinter S, Vrbova G. Delayed riluzole treatment is able to rescue injured rat spinal motoneurons. *Neuroscience*. 2007;144(2):431-438.
56. Stutzmann JM, Pratt J, Boraud T, Gross C. The effect of riluzole on post-traumatic spinal cord injury in the rat. *Neuroreport*. 1996;7(2):387-392.
57. Simard JM, Tsybalyuk O, Keledjian K, Ivanov A, Ivanova S, Gerzanich V. Comparative effects of glibenclamide and riluzole in a rat model of severe cervical spinal cord injury. *Exp Neurol*. 2012;233(1):566-574.
58. Schwartz G, Fehlings MG. Evaluation of the neuroprotective effects of sodium channel blockers after spinal cord injury: improved behavioral and neuroanatomical recovery with riluzole. *J Neurosurgery*. 2001;94(2 suppl):245-256.
59. Kwon BK, Roy J, Lee JH, et al. Magnesium chloride in a polyethylene glycol formulation as a neuroprotective therapy for acute spinal cord injury: preclinical refinement and optimization. *J Neurotrauma*. 2009;26(8):1379-1393.
60. Luo J, Borgens R, Shi R. Polyethylene glycol immediately repairs neuronal membranes and inhibits free radical production after acute spinal cord injury. *J Neurochem*. 2002;83(2):471-480.
61. Kaptanoglu E, Beskonakli E, Solaroglu I, Kilinc A, Taskin Y. Magnesium sulfate treatment in experimental spinal cord injury: emphasis on vascular changes and early clinical results. *Neurosurg Rev*. 2003;26(4):283-287.
62. Kaptanoglu E, Beskonakli E, Okutan O, Selcuk Surucu H, Taskin Y. Effect of magnesium sulphate in experimental spinal cord injury: evaluation with ultrastructural findings and early clinical results. *J Clin Neurosci*. 2003;10(3):329-334.
63. Kaptanoglu E, Beskonakli E, Solaroglu I, Kilinc A, Taskin Y. Magnesium sulfate treatment in experimental spinal cord injury: emphasis on vascular changes and early clinical results. *Neurosurg Rev*. 2003;26(4):283-287.
64. Chen M, Ona VO, Li M, et al. Minocycline inhibits caspase-1 and caspase-3 expression and delays mortality in a transgenic mouse model of Huntington disease. *Nat Med*. 2000;6(7):797-801.
65. Giuliani F, Fu SA, Metz LM, Yong VW. Effective combination of minocycline and interferon-beta in a model of multiple sclerosis. *J Neuroimmunol*. 2005;165(1-2):83-91.
66. Seabrook TJ, Jiang L, Maier M, Lemere CA. Minocycline affects microglia activation, Abeta deposition, and behavior in APP-tg mice. *Glia*. 2006;53(7):776-782.
67. Zanjani TM, Sabetkasaei M, Mosaffa N, Manaheji H, Labibi F, Farokhi B. Suppression of interleukin-6 by minocycline in a rat model of neuropathic pain. *Eur J Pharmacol*. 2006;538(1-3):66-72.
68. Lee SM, Yune TY, Kim SJ, et al. Minocycline inhibits apoptotic cell death via attenuation of TNF-alpha expression following iNOS/NO induction by lipopolysaccharide in neuron/glia co-cultures. *J Neurochem*. 2004;91(3):568-578.
69. Drabek T, Janata A, Wilson CD, et al. Minocycline attenuates brain tissue levels of TNF-alpha produced by neurons after prolonged hypothermic cardiac arrest in rats. *Resuscitation*. 2014;85(2):284-291.
70. Wells JE, Hurlbert RJ, Fehlings MG, Yong VW. Neuroprotection by minocycline facilitates significant recovery from spinal cord injury in mice. *Brain*. 2003;126(pt 7):1628-1637.
71. Festoff BW, Ameenuddin S, Arnold PM, Wong A, Santacruz KS, Citron BA. Minocycline neuroprotects, reduces microgliosis, and inhibits caspase protease expression early after spinal cord injury. *J Neurochem*. 2006;97(5):1314-1326.
72. Casha S, Zygun D, McGowan MD, Bains I, Yong VW, Hurlbert RJ. Results of a phase II placebo-controlled randomized trial of minocycline in acute spinal cord injury. *Brain*. 2012;135(pt 4):1224-1236.
73. Imanaka T, Hukuda S, Maeda T. The role of GM1-ganglioside in the injured spinal cord of rats: an immunohistochemical study using GM1-antisera. *J Neurotrauma*. 1996;13(3):163-170.
74. Geisler FD, Dorsey FC, Coleman William. Recovery of motor function after spinal-cord injury — a randomized, placebo-controlled trial with gm-1 ganglioside — NEJM. *N Engl J Med*. 1991;324(26):1829-1838.
75. Geisler FH, Coleman WP, Grieco G, Poonian D. The Sygen multicenter acute spinal cord injury study. *Spine*. 2001;26(24 suppl):S87-S98.
76. Siddiqui AM, Khazaei M, Fehlings MG. Translating mechanisms of neuroprotection, regeneration, and repair to treatment of spinal cord injury. *Prog Brain Res*. 2015;218:15-54.
77. Teng YD, Mocchetti I, Taveira-DaSilva AM, Gillis RA, Wrathall JR. Basic fibroblast growth factor increases long-term survival of spinal motor neurons and improves respiratory function after experimental spinal cord injury. *J Neurosci*. 1999;19(16):7037-7047.
78. Teng YD, Mocchetti I, Wrathall JR. Basic and acidic fibroblast growth factors protect spinal motor neurones in vivo after experimental spinal cord injury. *Eur J Neurosci*. 1998;10(2):798-802.
79. Wallner S, Peters S, Pitzer C, Resch H, Bogdahn U, Schneider A. The Granulocyte-colony stimulating factor has a dual role in neuronal and vascular plasticity. *Front Cell Dev Biol*. 2015;3:48.
80. Nishio Y, Koda M, Kamada T, et al. Granulocyte colony-stimulating factor attenuates neuronal death and promotes functional recovery after spinal cord injury in mice. *J Neuropathol Exp Neurol*. 2007;66(8):724-731.
81. Koda M, Nishio Y, Kamada T, et al. Granulocyte colony-stimulating factor (G-CSF) mobilizes bone marrow-derived cells into injured spinal cord and promotes functional recovery after compression-induced spinal cord injury in mice. *Brain Res*. 2007;1149:223-231.
82. Takahashi H, Yamazaki M, Okawa A, et al. Neuroprotective therapy using granulocyte colony-stimulating factor for acute spinal cord injury: a phase I/IIa clinical trial. *Eur Spine J*. 2012;21(12):2580-2587.
83. Kamiya K, Koda M, Furuya T, et al. Neuroprotective therapy with granulocyte colony-stimulating factor in acute spinal cord injury: a comparison with high-dose methylprednisolone as a historical control. *Eur Spine J*. 2015;24(5):963-967.
84. Kitamura K, Iwanami A, Fujiyoshi K, et al. Recombinant human hepatocyte growth factor promotes functional recovery after spinal cord injury. In: Uchida KNM, Ozawa H, Katoh S, Toyama Y, eds. *Neuroprotection and Regeneration of the Spinal Cord*. Japan: Springer; 2014:147-167.
85. Kitamura K, Fujiyoshi K, Yamane J, et al. Human hepatocyte growth factor promotes functional recovery in primates after spinal cord injury. *PLoS One*. 2011;6:e27706.

86. Kitamura K, Iwanami A, Nakamura M, et al. Hepatocyte growth factor promotes endogenous repair and functional recovery after spinal cord injury. *J Neurosci Res*. 2007;85(11):2332-2342.
87. Kwon BK, Mann C, Sohn HM, et al. Hypothermia for spinal cord injury. *Spine J*. 2008;8(6):859-874.
88. Dehaes M, Aggarwal A, Lin PY, et al. Cerebral oxygen metabolism in neonatal hypoxic ischemic encephalopathy during and after therapeutic hypothermia. *J Cereb Blood Flow Metab*. 2014;34(1):87-94.
89. Dingley J, Tooley J, Liu X, et al. Xenon ventilation during therapeutic hypothermia in neonatal encephalopathy: a feasibility study. *Pediatrics*. 2014;133(5):809-818.
90. Hypothermia after Cardiac Arrest Study Group. Mild therapeutic hypothermia to improve the neurologic outcome after cardiac arrest. *N Engl J Med*. 2002;346(8):549-556.
91. Lo TP Jr, Cho KS, Garg MS, et al. Systemic hypothermia improves histological and functional outcome after cervical spinal cord contusion in rats. *J Comp Neurol*. 2009;514(5):433-448.
92. Levi AD, Green BA, Wang MY, et al. Clinical application of modest hypothermia after spinal cord injury. *J Neurotrauma*. 2009;26(3):407-415.
93. Kwon BK, Curt A, Belanger LM, et al. Intrathecal pressure monitoring and cerebrospinal fluid drainage in acute spinal cord injury: a prospective randomized trial. *Journal Neurosurg Spine*. 2009;10(3):181-193.
94. Martirosyan NL, Kalani MY, Bichard WD, et al. Cerebrospinal fluid drainage and induced hypertension improve spinal cord perfusion after acute spinal cord injury in pigs. *Neurosurgery*. 2015;76(4):461-468; discussion 468-469.
95. Forgione N, Fehlings MG. Rho-ROCK inhibition in the treatment of spinal cord injury. *World Neurosurg*. 2014;82(3-4):e535-e539.
96. Fehlings MG, Theodore N, Harrop J, et al. A phase I/IIa clinical trial of a recombinant Rho protein antagonist in acute spinal cord injury. *J Neurotrauma*. 2011;28(5):787-796.
97. Bregman BS, Kunkel-Bagden E, Schnell L, Dai HN, Gao D, Schwab ME. Recovery from spinal cord injury mediated by antibodies to neurite growth inhibitors. *Nature*. 1995;378(6556):498-501.
98. Hamid S, Hayek R. Role of electrical stimulation for rehabilitation and regeneration after spinal cord injury: an overview. *Eur Spine J*. 2008;17(9):1256-1269.
99. Angeli CA, Edgerton VR, Gerasimenko YP, Harkema SJ. Altering spinal cord excitability enables voluntary movements after chronic complete paralysis in humans. *Brain*. 2014;137(pt 5):1394-1409.
100. Arriola A, Kiel ME, Shi Y, McKinnon RD. Adjunctive MSCs enhance myelin formation by xenogenic oligodendrocyte precursors transplanted in the retina. *Cell Res*. 2010;20(6):728-731.
101. Wang L, Shi J, van Ginkel FW, et al. Neural stem/progenitor cells modulate immune responses by suppressing T lymphocytes with nitric oxide and prostaglandin E2. *Exp Neurol*. 2009;216(1):177-183.
102. Okamura RM, Lebkowski J, Au M, Priest CA, Denham J, Majumdar AS. Immunological properties of human embryonic stem cell-derived oligodendrocyte progenitor cells. *J Neuroimmunol*. 2007;192(1-2):134-144.
103. Shi Y, Despons C, Do JT, Hahm HS, Scholer HR, Ding S. Induction of pluripotent stem cells from mouse embryonic fibroblasts by Oct4 and Klf4 with small-molecule compounds. *Cell Stem Cell*. 2008;3(5):568-574.
104. Salewski RP, Mitchell RA, Li L, et al. Transplantation of induced pluripotent stem cell-derived neural stem cells mediate functional recovery following thoracic spinal cord injury through remyelination of axons. *Stem Cells Transl Med*. 2015;4(7):743-754.
105. Ahuja C, Fehlings M. Bridging the gap: novel neuroregenerative and neuroprotective strategies in spinal cord injury. *Stem Cells Transl Med*. 2016 5(7):914-924.
106. Dasari VR, Veeravalli KK, Dinh DH. Mesenchymal stem cells in the treatment of spinal cord injuries: A review. *World J Stem Cells*. 2014;6(2):120-133.
107. Quertainmont R, Cantinieux D, Botman O, Sid S, Schoenen J, Franzen R. Mesenchymal stem cell graft improves recovery after spinal cord injury in adult rats through neurotrophic and pro-angiogenic actions. *PLoS One*. 2012;7(6):e39500.
108. Kim JW, Ha KY, Molon JN, Kim YH. Bone marrow-derived mesenchymal stem cell transplantation for chronic spinal cord injury in rats: comparative study between intrathecal and intravenous transplantation. *Spine*. 2013;38(17):E1065-E1074.
109. Swartzlander MD, Blakney AK, Amer LD, Hankenson KD, Kyriakides TR, Bryant SJ. Immunomodulation by mesenchymal stem cells combats the foreign body response to cell-laden synthetic hydrogels. *Biomaterials*. 2015;41:79-88.
110. Bessout R, Semont A, Demarquay C, Charcosset A, Benderitter M, Mathieu N. Mesenchymal stem cell therapy induces glucocorticoid synthesis in colonic mucosa and suppresses radiation-activated T cells: new insights into MSC immunomodulation. *Mucosal Immunol*. 2014;7(3):656-669.
111. Lim JH, Kim JS, Yoon IH, et al. Immunomodulation of delayed-type hypersensitivity responses by mesenchymal stem cells is associated with bystander T cell apoptosis in the draining lymph node. *J Immunol*. 2010;185(7):4022-4029.
112. Williams RR, Bunge MB. Schwann cell transplantation: a repair strategy for spinal cord injury? *Prog Brain Res*. 2012;201:295-312.
113. Windus LC, Lineburg KE, Scott SE, et al. Lamellipodia mediate the heterogeneity of central olfactory ensheathing cell interactions. *Cell Mol Life Sci*. 2010;67(10):1735-1750.
114. Silva NA, Cooke MJ, Tam RY, et al. The effects of peptide modified gellan gum and olfactory ensheathing glia cells on neural stem/progenitor cell fate. *Biomaterials*. 2012;33(27):6345-6354.
115. Zhang J, Chen H, Duan Z, et al. The effects of co-transplantation of olfactory ensheathing cells and schwann cells on local inflammation environment in the contused spinal cord of rats. *Mol Neurobiol*. 2016.
116. Ekberg JA, St John JA. Olfactory ensheathing cells for spinal cord repair: crucial differences between subpopulations of the glia. *Neural Regen Res*. 2015;10(9):1395-1396.
117. Ibrahim AG, Kirkwood PA, Raisman G, Li Y. Restoration of hand function in a rat model of repair of brachial plexus injury. *Brain*. 2009;132(pt 5):1268-1276.
118. Liu J, Chen P, Wang Q, et al. Meta analysis of olfactory ensheathing cell transplantation promoting functional recovery of motor nerves in rats with complete spinal cord transection. *Neural Regen Res*. 2014;9(20):1850-1858.
119. Li L, Adnan H, Xu B, et al. Effects of transplantation of olfactory ensheathing cells in chronic spinal cord injury: a systematic review and meta-analysis. *Eur Spine J*. 2015;24(5):919-930.
120. Caicco MJ, Zahir T, Mothe AJ, et al. Characterization of hyaluronan-methylcellulose hydrogels for cell delivery to the injured spinal cord. *J Biomed Mater Res A*. 2013;101(5):1472-1477.
121. Mothe AJ, Tam RY, Zahir T, Tator CH, Shoichet MS. Repair of the injured spinal cord by transplantation of neural stem cells in a hyaluronan-based hydrogel. *Biomaterials*. 2013;34(15):3775-3783.
122. Tam RY, Cooke MJ, Shoichet MS. A covalently modified hydrogel blend of hyaluronan-methyl cellulose with peptides and growth factors influences neural stem/progenitor cell fate. *J Mater Chem*. 2012;22(37):19402-19411.
123. Ansorena E, De Berdt P, Ucakar B, et al. Injectable alginate hydrogel loaded with GDNF promotes functional recovery in a hemisection model of spinal cord injury. *Int J Pharm*. 2013;455(1-2):148-158.
124. Itosaka H, Kuroda S, Shichinohe H, et al. Fibrin matrix provides a suitable scaffold for bone marrow stromal cells transplanted into injured spinal cord: a novel material for CNS tissue engineering. *Neuropathology*. 2009;29(3):248-257.
125. Zweckberger K, Ahuja CS, Liu Y, Wang J, Fehlings MG. Self-assembling peptides optimize the post-traumatic milieu and synergistically enhance the effects of neural stem cell therapy after cervical spinal cord injury. *Acta Biomater*. 2016;42:77-89.
126. Kumar R, DiMenna L, Schrode N, et al. AID stabilizes stem-cell phenotype by removing epigenetic memory of pluripotency genes. *Nature*. 2013;500(7460):89-92.
127. Lu P, Wang Y, Graham L, et al. Long-distance growth and connectivity of neural stem cells after severe spinal cord injury. *Cell*. 2012;150(6):1264-1273.
128. Bracken MB. Methylprednisolone in the management of acute spinal cord injuries. *Med J Aust*. 1990;153(6):368.
129. Bracken MB, Collins WF, Freeman DF, et al. Efficacy of methylprednisolone in acute spinal cord injury. *JAMA*. 1984;251(1):45-52.
130. Martin AR, Cohen-Adad J, Tarmohamed Z, et al. Translating state-of-the-art spinal cord MRI techniques to clinical use: a systematic review of clinical studies utilizing DTI, MT, MWF, MRS, and fMRI. *Neuroimage Clin*. 2015;10:192-238.
131. Cadotte DW, Fehlings MG. Will imaging biomarkers transform spinal cord injury trials? *Lancet Neurol*. Sep 2013;12(9):843-844.

132. Kwon BK, Streijger F, Fallah N, et al. Cerebrospinal fluid biomarkers to stratify injury severity and predict outcome in human traumatic spinal cord injury. *J Neurotrauma*. Feb 2017;34(3):567-580.
133. Kwon BK, Stammers AM, Belanger LM, et al. Cerebrospinal fluid inflammatory cytokines and biomarkers of injury severity in acute human spinal cord injury. *J Neurotrauma*. 2010;27(4):669-682.
134. Nieto-Diaz M, Esteban FJ, Reigada D, et al. MicroRNA dysregulation in spinal cord injury: causes, consequences and therapeutics. *Front Cell Neurosci*. 2014;8:53.
135. Martin AR, Aleksanderek I, Fehlings MG. Diagnosis and acute management of spinal cord injury: current best practices and emerging therapies. *Curr Trauma Rep*. 2015;1(3):169-181.
136. Kirshblum SC, Waring W, Biering-Sorensen F, et al. Reference for the 2011 revision of the International Standards for Neurological Classification of Spinal Cord Injury. *J Spinal Cord Med*. 2011;34(6):547-554.
137. Singh A, Tetreault L, Kalsi-Ryan S, Nouri A, Fehlings MG. Global prevalence and incidence of traumatic spinal cord injury. *Clin Epidemiol*. 2014;6:309-331.
138. Ahuja C, Fehlings M. How can we regenerate the traumatically injured spinal cord?, 2016. <http://www.neurology-central.com/2016/06/01/how-can-we-regenerate-the-traumatically-injured-spinal-cord/>. Accessed August 8, 2016.

Acknowledgments

We thank Madeleine O'Higgins for copyediting and Chi Lam for organizational support.